



Colombian Journal of Anesthesiology

Revista Colombiana de Anestesiología

www.revcolanest.com.co

OPEN

Wolters Kluwer

How many people need palliative care for cancer and non-cancer diseases in a middle-income country? Analysis of mortality data

¿Cuántas personas requieren cuidados paliativos para enfermedades oncológicas y no oncológicas en un país de ingresos medios? Análisis de datos de mortalidad

Jose A. Calvache^a, Fabian Gil^b, Esther de Vries^b

^a Department of Anesthesiology, Epidemiology and Biostatistics Center, Universidad del Cauca, Popayan, Colombia

^b PhD Program in Clinical Epidemiology, Department of Clinical Epidemiology and Biostatistics, Pontificia Universidad Javeriana, Bogotá, Colombia.

Keywords: Palliative care, Needs assessment, Public health, Health services needs and demand, Delivery of health care, End-of-life care, Terminal care

Palabras clave: Cuidados paliativos, Evaluación de las necesidades, Salud pública, Necesidades de servicios de salud y demanda, Atención en salud, Cuidado al final de la vida, Cuidado terminal

Abstract

Introduction: Aging of the population and the accompanying increase in prevalence of chronic illnesses mean that more people will need palliative care. This need has not been extensively quantified in middle-income countries to policy planning and expanding health care.

Objective: Provide an estimate of the need of palliative care services in Colombia and compare these needs with the current available offer.

Methods: Cross-sectional study based on mortality statistics for Colombia for the period 2012 to 2016. We calculated age-specific and sex-specific numbers of deaths and mortality rates from death certificates for defined chronic illnesses to estimate the prevalence of palliative care need in Colombia, and contrasted

this information with the current offer according to the Colombian Observatory of Palliative Care.

Results: The numbers of deaths requiring palliative care increased from 107,065 in 2012 to 128,670 in 2016 (61.2% of total deaths). The causes of these deaths vary by age group, with a clearly more important proportion of heart and cerebrovascular diseases and dementia in advanced ages, and HIV/AIDS in young ages. In all age groups, malignant neoplasms are an important part of the causes of deaths of those requiring palliative care (31.3% of all deaths in 2016). Contrasting these needs, in most areas there is no or very limited offer of palliative care services.

Conclusion: A real palliative care policy, including a vast increase in training opportunities in the field and regulation allowing a wide range of health practitioners to be involved in palliative care, is necessary in Colombia to improve the palliative care offer.

How to cite this article: Calvache JA, Gil F, de Vries E. How many people need palliative care for cancer and non-cancer diseases in a middle-income country? Analysis of mortality data. Colombian Journal of Anesthesiology. 2020;00:000-000.

Read the Spanish version of this article at: <http://links.lww.com/RCA/A928>

Copyright © 2020 Sociedad Colombiana de Anestesiología y Reanimación (S.C.A.R.E.). Published by Wolters Kluwer. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Correspondence: Department of Clinical Epidemiology and Biostatistics, Pontificia Universidad Javeriana, Hospital San Ignacio, Piso 2, Cra. 7 No 40-62, Bogotá, Colombia. E-mail: estherdevries@javeriana.edu.co

Colombian Journal of Anesthesiology (2020) Vol:No

<http://dx.doi.org/10.1097/CJ9.0000000000000159>

Resumen

Introducción: El envejecimiento de la población y el consecuente aumento en la prevalencia de enfermedades crónicas implica que habrá más personas que necesitarán cuidados paliativos. Esta necesidad no se ha cuantificado mayormente en los países de medianos ingresos con miras a la planificación de políticas y a la ampliación de la atención en salud.

Objetivo: Ofrecer un estimado de la necesidad de servicios de cuidados paliativos en Colombia y comparar dichas necesidades con la oferta disponible en la actualidad.

Métodos: Estudio transversal basado en estadísticas de mortalidad en Colombia para el periodo comprendido entre 2012–2016. Calculamos las cifras de muertes y tasas de mortalidad específicas por edad y sexo a partir de los certificados de defunción correspondientes a enfermedades crónicas definidas, a fin de estimar la prevalencia de la necesidad de cuidados paliativos en Colombia, y se comparó dicha información con la oferta actual, de acuerdo con el Observatorio Colombiano de Cuidados Paliativos.

Resultados: Las cifras correspondientes a muertes que requirieron cuidados paliativos aumentaron de 107 065 en el 2012 a 128 670 en el 2016 (61.2% del total de muertes). Las causas de estas muertes varían según el grupo etario, con un porcentaje evidentemente mayor de enfermedades cardíacas, cerebrovasculares y demencia en edades avanzadas, y VIH/SIDA en los grupos más jóvenes. En todos los grupos etarios, las neoplasias malignas son una parte importante de las causas de muerte de quienes requieren cuidados paliativos (31.3% de todas las muertes en el 2016). Estas necesidades contrastan con el hecho de que en la mayoría de las áreas no existen servicios de cuidados paliativos, o su oferta es muy limitada.

Conclusiones: En Colombia se necesita una verdadera política de cuidados paliativos, incluyendo un aumento significativo de las oportunidades de capacitación en el área, así como un marco regulatorio que permita que una amplia gama de profesionales de la salud participen en cuidados paliativos, a fin de mejorar la oferta de estos servicios.

What is already known about the topic?

- People with a wide range of chronic conditions get benefit from palliative care at the end of life.
- Estimates from routine mortality data can give rough but reliable estimates of the population's need for palliative care.

What this paper adds?

- The proportion of individuals in need of palliative care at the end of life varies significantly in Colombia and is increasing over time.
- The current offer of palliative care services in Colombia seriously lags behind the current demand.
- There is an urgent need of effective policy to increase the capacity of palliative care services across the country, especially in less-populated areas.

Introduction

The need for palliative health care has been defined as “the population’s ability to benefit from palliative care.”^{1,2} This designation includes a professional perspective, which is useful when considering the need for palliative care at a national population level.³ In epidemiology, “need” refers to “specific indicators of disease which require intervention because the level is above that generally accepted within a particular society.”⁴ Following this definition, “needs” must reflect value judgments on the ability to control the problem but also awareness about certain growing problems within societies.⁵

Demographic changes leading to aging of the population and the accompanying increase in long-term conditions and multimorbidity requiring advanced medical care have moved all health sectors to provide day-to-day care to patients with more advanced diseases and end-of-life. Many people, not only terminally ill patients, could benefit from palliative care which is mainly focused on quality of life and relief of symptoms.^{6,7}

It is well known that worldwide, there is a big discrepancy in availability of palliative care services between countries, with most low-income and middle-income countries having clearly unmet needs. In this study, we set out to determine the needs of palliative care for cancer and non-cancer diseases in Colombia, which is considered an upper middle-income country by the World Bank but as well a country with many challenges in society and very large socioeconomic inequalities between and within regions of the country.^{8,9}

The demographic composition of the Colombian population is changing rapidly, due to general demographic changes, combined with a large influx of refugees from Venezuela and the recent “peace” process, which has greatly diminished deaths due to violence. In general, the tendency is for the population ages 65+ to proportionally increase rapidly—which almost automatically implies strong increases in the prevalence of non-communicable diseases.¹⁰ This translates into an expected, but currently undocumented, increased need for palliative care. Understanding healthcare needs is essential to planning and expanding services, yet no national palliative care needs estimation has been carried out in Colombia nor, as far as we know, in South America.

Since 2014, the Colombian government has introduced regulation regarding health care of people with advanced, degenerative, terminal, and irreversible diseases.¹¹ In addition, a guideline for the treatment of people in palliative care is available, as well as access to opioids for pain management.^{12,13}

Patients with a large variety of conditions are in need of palliative care. A particular, and frequently studied group of diseases is cancer—which need many different types of palliative support, from nutrition to ventilation and from pain management to psychologic and spiritual counsel-

ing.^{14–16} However, many other diseases are also very important in the need for palliative care.^{17,18} Therefore, in this project we estimate the potential population in need of palliative care services in Colombia based on national mortality data—disaggregated by cancer and non-cancer deaths. Using the available data from the national palliative care observatory, we intend to compare the estimated needs with the current offer, to help decision makers in determining where the greatest needs are.

Provide an estimate of the need of palliative care services in Colombia and compare these needs with the current available offer.

Methods

We developed an observational cross-sectional death certificate-based study of individuals deceased above 18 years in Colombia between 2012 and 2016. The analyses are based on public and anonymized data from death certificates provided by National Administrative Department of Statistics of Colombia (DANE). Vital records are continuous statistics that collect information on births, fetal and non-fetal deaths, thus providing the information on changes in the levels and patterns of mortality and fertility. These information is publicly available through website of DANE (<https://www.dane.gov.co>). Chronic conditions for which palliative care is required were identified from the main cause of death using ICD-10 codes.

Deaths requiring palliative care

A minimal estimate was calculated based on the number of deaths from specific conditions identified as having palliative care needs.³ Unfortunately, it was not possible to link with hospital records, limiting the possibility of more refined estimates. In line with the methods applied by Gomes et al,¹⁹ in their modification to a previous paper,³ we defined deaths requiring palliative care as those deaths with underlying cause of death coded as summarized in Table 1. These deaths-related codes have shown a requirement of palliative care ranging from 69% to 82%.³

Offer of palliative care services

Data to estimate the current offer of palliative care services was extracted from the Colombian Observatory of Palliative Care, which provides information about availability of palliative care services per 100,000 inhabitants on a departmental level. These estimates are based on current hospitality offer of services related to palliative care and registered as health providers including day care, ambulatory palliative medical consultations, short or long stay hospital-based palliative care, residential-based palliative care, and home-based palliative care.¹⁷

Table 1. Description of Cause of Death and ICD-10 codes used to categorize deaths requiring palliative care.

Underlying cause of death	ICD-10 codes
Malignant neoplasm	C00–C97
Heart & cerebrovascular disease	I00–I52 (excluding I12 and I13.1), I60–I69
Renal disease	N17, N18, N28, I12, I13.1
Liver disease	K70–K77
Respiratory disease	J06–J18, J20–J22, J40–J47, J96
Neurodegenerative disease	G10, G20, G35, G122, G903, G231
Alzheimer's, dementia and senility	F01, F03, G30, R54
HIV/AIDS	B20–B24

Source: Authors.

Analysis

We stratified the analyses for cancer and other conditions (as specified in Table 1), age groups and major cities. We considered as variables of interest the total population (and stratified by age) and the total (and per category) number of deaths. R software (version 3.5.0, R 3.5.0: R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2019. Disponible en: <https://www.R-project.org/>) was used for data analysis and graphs.

Ethical approval

Since this study was classified as a study without risk, based on routinely, anonymized and publicly available data, no ethical approval or informed consent was necessary for this study.

Results

Between 2012 and 2016, the Colombian estimated population grew from 46,581,823 to 48,747,708 persons. In the same period, the proportion of the population aged 60 years and over increased from 10.3% to 11.4% of the total population.

Figure 1 illustrates the increase in numbers of deaths due to any cause as well as those due to diseases which likely require palliative care: the absolute numbers of deaths increased from 185,248 in 2012 to 210,224 in 2016; the numbers of deaths requiring palliative care according to our definition increased from 107,065 in 2012 to 128,670 in 2016 (61.2%). About 1/3 of these cases were cancer-related (31.3% in 2016).




Figure 1. Absolute number of deaths due to any cause and proportion of deaths potentially requiring palliative care from 2012 to 2016 in Colombia. Source: Authors.

The number of deaths likely to require palliative care increased with age, over the study period there were 7667 deaths requiring palliative care in the age group 18 to 44 years, increasing to 26,840 for ages 45 to 64, 42,288 for 65 to

79 and 48,203 for the 80+ population. Figure 2 shows how the causes of these deaths vary by age group, with a clearly more important proportion of heart and cerebrovascular diseases as well as dementia in advanced age-groups, and




Figure 2. Number of deaths by age groups and causes from 2012 to 2016 in Colombia. Source: Authors.




Figure 3. (A) People potentially requiring palliative care per 100,000 person-years. (B) Offer of palliative care services in Colombia. Source: Authors.

HIV/AIDS being much more important in the younger segments of the population. In all age groups, malignant neoplasms are an important part of the causes of deaths of those requiring palliative care.

Figure 3A shows the people potentially requiring palliative care per 100,000 person-years—with concentrations in the more densely populated areas of the country. Contrasting this need of palliative care with the formal offer of palliative care services as reported by the national observatory of palliative care (Fig. 3B), it is clear that in some areas, there is no offer at all, in other areas with high need, the offer is limited, whereas in some areas with high demand the offer is below national average.

Discussion

Our results clearly show the high and growing need for palliative care services in the population, and also the deficiency of palliative care services on offer. Estimates from routine mortality data can give rough but reliable estimates of the population's need for palliative care.^{3,18,19} This information is essential to plan services. A real palliative care policy, not only on paper but implemented in reality, is necessary in Colombia to improve the palliative care service offer. Our findings clearly indicate a high potential need for palliative care in many areas of the country where this type of care is not currently widely offered.

More palliative care support is expected to be required for some illnesses and clinical syndromes, such as patients with multimorbidity, chronic progressive illnesses with long disease courses, diseases with complex symptoms and demographic changes. Our estimates show

that the proportion of total deaths potentially requiring palliative care increases with time and they are in agreement with results from other middle-income and high-income countries.^{3,7,20} The proportion of individuals who died from diseases that indicate palliative care needs at the end of life ranged in the literature from 38% to 74% with an important cross-country variation.^{20–22} At population level, these estimations provide robust indications of the challenge countries are facing without major differences between high-income and middle-income countries.²⁰ Our estimates of palliative care needs are probably quite similar to those of other Latin-American countries in transition, the palliative care offer depends heavily on the health systems in place in each country.

It is as yet very unclear how exactly health professionals and the entire Colombian health care system are to meet the increased population palliative care need. The current offer of palliative care is mainly due to services in third level of care (high complexity services) with 13 programs in the country, only 1 program at a second level hospital, 3 multi-level services and around 60 palliative home care services for the whole country (low-complexity services).²³ Despite the strong increase between 2012 and 2016 in the offer of palliative care services (an increase of 500%), those services are located in urban big cities and departments (i.e., Bogotá D.C., Antioquia, Valle del Cauca, and Atlántico). Unfortunately, there were no data available regarding the capacity and scope of those providers at national level. Our data show that there are several areas with no offer at all in terms of palliative care.

There is a clear and urgent need for training in palliative care in medical faculties and adaptations from the side of the health system and health insurers to facilitate all kind

of palliative care services. The current number of palliative care specialists is still very low and required early training is scarce. Research from low-income and middle-income countries highlights the need for education and training of staff and healthcare practitioners in palliative care and palliative care services.²⁴ Besides training in palliative care skills at all levels, legal reforms will be necessary as the current legislation leaves little to no room for a wide range of health practitioners to be involved, to prescribe or apply, for example, morphine derivatives or nutritional interventions. In 2012, there were 57 medical faculties in Colombia and only 3 included palliative care training as an integral part of their health sciences curriculum.²³ All these faculties and current palliative care services are located in urban big cities without major covering of middle cities or rural areas.

Limitations

Of course, a study based on death certificate information suffers from all factors that influence the quality of the death certification. Fortunately, in the studied period, a very large proportion of the deaths included in our analyses were certified by medical doctors, cause of death coding is always tricky.²⁵ Some patients suffering from the diseases under study may have died of other diseases and not have been included. In some remote areas and particularly in isolated populations, such as some indigenous tribes, not all deaths may have been reported. However, we believe that by and large, our estimates reflect the need of palliative care—in all of its forms—in Colombia in recent years. Considering that not only fatally ill people need palliative care for optimum treatment, our estimates are likely a conservative estimate—the real need may be substantially higher. The contrast with the current official offer of palliative care is stark. Strong legislation regarding opioid availability is limiting availability of vital pain management medicines,²⁶ and lack of knowledge and human resources for all aspects of palliative care is clearly lacking.

Conclusion

The current offer of palliative care services seriously lags behind the current demand. With very limited training facilities in the field of palliative care and a growing need because of the rapidly ageing populations, effective policy is urgently needed to improve the situation.

Ethical responsibilities

Design. Observational cross-sectional death-certificate based study of individuals deceased above 18 years in Colombia between 2012 and 2016.

Data sources. The analyses are based on public and anonymized data from death certificates provided by

National Administrative Department of Statistics of Colombia (DANE). Chronic conditions for which palliative care is required were identified from the main cause of death using ICD-10 codes.

Ethical approval. Since this study used routinely, anonymized, and publicly available data, no ethical approval was necessary for this study.

Acknowledgments

We would like to thank DANE and all officials signing the death certificates and checking the coding for their dedication and work; without them this study would not have been possible.

Authorship

All authors have made substantive intellectual contributions to the development of this article in terms of conceptualized the study, data extraction and analysis, supervision of the article preparation and writing of the final version.

Funding

The author(s) disclosed receipt of the following financial support. This project was funded by a grant from Colciencias 777-2017 Convocatoria para proyectos de ciencia, tecnología e innovación en Salud 2017. Project name “Decisiones medicas al final de la vida en pacientes oncológicos en Colombia.”

Conflicts of interest

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

References

1. Culyer AJ. Need: the idea won't do – but we still need it. *Soc Sci Med* 1995;40:727–730.
2. Bradshaw J. A taxonomy of social need. *New Soc* 1972;30:640–643.
3. Murtagh FE, Bausewein C, Verne J, et al. How many people need palliative care? A study developing and comparing methods for population-based estimates. *Palliat Med* 2014;28:49–58.
4. Porta M. *A Dictionary of Epidemiology*. 6th ed. Oxford: Oxford University Press; 2014.
5. Franks PJ, Salisbury C, Bosanquet N, et al. The level of need for palliative care: a systematic review of the literature. *Palliat Med* 2000;14:93–104.
6. World Health Organization. WHO Definition of Palliative Care. 2011; World Health Organization, Geneva: <http://www.who.int/cancer/palliative/definition/en/>. [Accessed 16 January, 2019]
7. Etkind SN, Bone AE, Gomes B, et al. How many people will need palliative care in 2040? Past trends, future projections and implications for services. *BMC Med* 2017;15:102.
8. World Bank. 2018. Atlas de los Objetivos de Desarrollo Sostenible 2018 : Basado en los Indicadores del Desarrollo Mundial. World

- Bank Atlas; Washington, DC: World Bank. © World Bank. <https://openknowledge.worldbank.org/handle/10986/29788> License: CC BY 3.0 IGO.
9. Alvarado F, Chancel L, Piketty T, Saez E, Zucman G, editores. World inequality report 2018. Cambridge, Massachusetts: The Belknap Press of Harvard University Press; 2018. p. 332. <https://wir2018.wid.world/files/download/wir2018-full-report-english.pdf>. [Accessed 20 April, 2019].
 10. Demographic Observatory of Latin America. Demographic Observatory of Latin America 2016: Population Projections. 2016; Economic Commission for Latin America and the Caribbean (ECLAC), Demographic Observatory, Santiago:(LC/PUB.2017/3-P), Santiago, 2016. <http://www.cepal.org/es/publicaciones/41018-observatorio-demografico-america-latina-2017-proyecciones-poblacion-demographic>. [Accessed 20 April, 2019]
 11. Congreso de la República de Colombia. Diario Oficial No. 49.268 de 8 de septiembre de 2014. http://www.secretariasenado.gov.co/senado/basedoc/ley_1733_2014.html. [2014, Accessed 20 May, 2019].
 12. Ministerio de SaludGuía de práctica clínica Cuidados paliativos Adopción 2016. Ministerio de Salud y Protección Social; 2016; http://gpc.minsalud.gov.co/gpc_sites/Repositorio/Otros_conv/GPC_paliativo/gpc_paliativo.aspx [Accessed 30 May, 2019]
 13. Ministerio de Salud y Protección SocialCircular No. 022 de 2016 Lineamientos y Directrices para la gestión del acceso a medicamentos opioides para el manejo del dolor. Ministerio de Salud y Protección Social; 2016; <https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/DE/DIJ/circular-022-2016.pdf>. [Accessed 30 May, 2019]
 14. Bausewein C, Simon ST, Pralong A, et al. Palliative care of adult patients with cancer. *Dtsch Arztebl Int* 2015;112:863–870.
 15. Lorenz KA, Lynn J, Dy SM, et al. Evidence for improving palliative care at the end of life: a systematic review. *Ann Intern Med* 2008;148:147–159.
 16. Pergolizzi JV, Gharibo C, Ho K. Treatment considerations for cancer pain: a global perspective. *Pain Pract* 2015;15:778–792.
 17. Universidad El BosqueAnuario del Observatorio Colombiano De Cuidados Paliativos 2016. Universidad del Bosque; 2016; <http://occp.com.co/wp-content/themes/occp/docs/anuario-occp-2016.pdf>. [Accessed 27 June, 2019]
 18. Gómez-Batiste X, Martínez-Muñoz M, Blay C, et al. Prevalence and characteristics of patients with advanced chronic conditions in need of palliative care in the general population: a cross-sectional study. *Palliat Med* 2014;28:302–311.
 19. Gomes B, Pinheiro MJ, Lopes S, et al. Risk factors for hospital death in conditions needing palliative care: nationwide population-based death certificate study. *Palliat Med* 2018;32:891–901.
 20. Morin L, Aubry R, Frova L, et al. Estimating the need for palliative care at the population level: a cross-national study in 12 countries. *Palliat Med* 2016;31:526–536.
 21. Kane PM, Daveson BA, Ryan K, et al. The need for palliative care in Ireland: a population-based estimate of palliative care using routine mortality data, inclusive of nonmalignant conditions. *J Pain Symptom Manage* 2015;49:726–733.e1.
 22. Rosenwax LK, McNamara B, Blackmore AM, et al. Estimating the size of a potential palliative care population. *Palliat Med* 2005;19:556–562.
 23. Pastrana T, De Lima L, Wenk R, et al. Atlas de Cuidados Paliativos de Latinoamérica ALCP. 1st ed.Houston: IAHP Press; 2012.
 24. Maharaj S, Harding R. The needs, models of care, interventions and outcomes of palliative care in the Caribbean: a systematic review of the evidence. *BMC Palliat Care* 2016;15:9.
 25. Harteloh P, de Bruin K, Kardaun J. The reliability of cause-of-death coding in The Netherlands. *Eur J Epidemiol* 2010;25:531–538.
 26. Leon MX, De Lima L, Florez S, et al. Improving availability of and access to opioids in Colombia: description and preliminary results of an action plan for the country. *J Pain Symptom Manage* 2009;38:758–766.